

Policy Title:	Ilumya (tildrakizumab-asmn) subcutaneous		
		Department:	PHA
Effective Date:	01/01/2020		
Review Date:	9/18/2019 , 12/20/2019		
Revision Date:	9/18/2019, 12/20/2019		

Purpose: To support safe, effective and appropriate use of Ilumya (tildrakizumab-asmn).

Scope: Medicaid, Exchange, Medicare-Medicaid Plan (MMP)

Policy Statement:

Ilumya (tildrakizumab-asmn) is covered under the Medical Benefit when used within the following guidelines. Use outside of these guidelines may result in non-payment unless approved under an exception process.

Procedure:

Coverage of Ilumya (tildrakizumab-asmn) will be reviewed prospectively via the prior authorization process based on criteria below.

Initial Criteria:

Moderate to severe plaque psoriasis

- Patient must be 18 years of age or older: AND
- Patient has a diagnosis of moderate to severe plaque psoriasis: AND
Patient has a pretreatment tuberculosis (TB) screening with a TB skin test or an interferon gamma release assay (e.g., QFT-GIT, T-SPOT.TB); AND
- Is prescribed by, or in consultation with, a specialist in dermatology or rheumatology; AND
- At least 10% of BSA is affected OR crucial body areas (i.e., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected; AND
- Member meets either of the following:
 - Member has had an inadequate response to at least a 3 month trial methotrexate, cyclosporine or acitretin, or experienced clinically significant adverse effects from methotrexate, cyclosporine or acitretin; OR
 - Member has had an inadequate response to at least a 3 month trial of phototherapy (e.g., UVB, PUVA), unless intolerance experienced; AND
- Ilumya will not be used concomitantly with any other biologic DMARD or targeted synthetic DMARD.

Continuation of Therapy Criteria:

- Authorization may be granted for all members (including new members) who meet all initial authorization criteria and achieve or maintain positive clinical response within FDA dosing guidelines after at least 4 months of therapy with Ilumya as evidenced by low disease activity or improvement in signs and symptoms of the condition.

Coverage durations:

- Initial coverage: 6 months
- Continuation of therapy coverage: 6 months

*** Requests will also be reviewed to National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) if applicable. ***

Dosage/Administration:

Indication	Dose	Maximum Dosing (1 billable unit = 1 mg)
Plaque Psoriasis	100 mg subcutaneously at Week 0 and 4 then 100 mg every 12 weeks thereafter. Ilumya should be administered by a health care provider only	<u>Loading:</u> 100 units (100 mg) at Week 0 & 4 <u>Maintenance:</u> 100 units (100 mg) every 12 weeks

Appendix:

Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine or Acitretin:

- Alcoholism, alcoholic liver disease or other chronic liver disease
- Breastfeeding
- Cannot be used due to risk of treatment-related toxicity
- Drug interaction
- Pregnancy or planning pregnancy (male or female)
- Significant comorbidity prohibits use of systemic agents (examples include liver or kidney disease, blood dyscrasias, uncontrolled hypertension)

Investigational use: All therapies are considered investigational when used at a dose or for a condition other than those that are recognized as medically accepted indications as defined in any one of the following standard reference compendia: American Hospital Formulary Service Drug information (AHFS-DI), Thomson Micromedex DrugDex, Clinical Pharmacology, Wolters Kluwer Lexi-Drugs, or Peer-reviewed published medical literature indicating that sufficient evidence exists to support use. Neighborhood does not provide coverage for drugs when used for investigational purposes.

Applicable Codes:

Below is a list of billing codes applicable for covered treatment options. The below tables are provided for reference purposes and may not be all inclusive. Requests received with codes from tables below do not guarantee coverage. Requests must meet all criteria provided in the procedure section.

The following HCPCS/CPT code is:

HCPCS/CP T Code	Description
J3245	Injection, tildrakizumab, 1 mg

References:

1. Ilumya [package insert]. Whitehouse Station, NJ: Merck & Co., Inc.; March 2018.
2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol*. 2009;61:451-485.
3. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011;65(1):137-174.