

SPECIALTY GUIDELINE MANAGEMENT

OTEZLA (apremilast)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

1. Moderate to severe plaque psoriasis
2. Active psoriatic arthritis
3. Oral ulcers associated with Behçet's disease

All other indications are considered experimental/investigational and are not a covered benefit.

II. CRITERIA FOR INITIAL APPROVAL

A. **Moderate to severe plaque psoriasis**

Authorization of 6 months may be granted for treatment of moderate to severe plaque psoriasis for members who are 18 years of age or older when all of the following criteria are met:

1. Otezla is prescribed by, or in consultation with, a specialist in dermatology or rheumatology.
2. At least 10% of BSA is affected OR crucial body areas (i.e., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
3. Member meets either of the following criteria:
 - a. Member has had an inadequate response to at least a 3 month trial of methotrexate, cyclosporine or acitretin, or experienced clinically significant adverse effects from methotrexate, cyclosporine or acitretin
 - b. Member has had an inadequate response to at least a 3 month trial of phototherapy (e.g., UVB, PUVA), unless intolerance experienced
4. Otezla will not be used concomitantly with any other biologic DMARD (e.g. adalimumab, infliximab).

B. **Active psoriatic arthritis (PsA)**

Authorization of 6 months may be granted for treatment of active psoriatic arthritis (PsA) for members who are 18 years of age or older when all of the following criteria are met:

1. Otezla is prescribed by, or in consultation with, a specialist in dermatology or rheumatology.
2. Documented moderate to severe active disease and member meets either of the following criteria:
 - a. If member has predominantly axial disease or active enthesitis and/or dactylitis, member has experienced an inadequate response or intolerance to at least two non-steroidal anti-inflammatory drugs (NSAIDs), unless use is contraindicated
 - b. If member has peripheral arthritis, member has experienced an inadequate response to at least a 3 month trial of one oral disease-modifying anti-rheumatic agent (DMARD) such as methotrexate, azathioprine, sulfasalazine, or hydroxychloroquine

C. Behcet's syndrome

Authorization of 6 months may be granted for the treatment of oral ulcers associated with Behcet's syndrome when the member has had an inadequate response to at least one nonbiologic medication for Behcet's disease (e.g., colchicine, systemic glucocorticoids, azathioprine).

III. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for all members (including new members) who achieve or maintain positive clinical response after at least 4 months of therapy with Otezla as evidenced by low disease activity or improvement in signs and symptoms of the condition.

IV. QUANTITY LIMIT

1. Otezla 30mg tablet – 2 tablets per day
2. Otezla Starter Therapy Pack – 55 tablets per 28 days

V. REFERENCES

1. Otezla [package insert]. Summit, NJ: Celgene Corporation; July 2019.
2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol*. 2009;61:451-485.
3. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011;65(1):137-174.
4. Coates LC, Kavanaugh A, Mease PJ, et al. Group for research and assessment of psoriasis and psoriatic arthritis 2015 treatment recommendation for psoriatic arthritis. *Arthritis Rheumatol*. 2016 May;68(5):1060-71.