

Effective Date: 12/01/2020
Last Reviewed: 9/2020
Scope: Medicaid

Ofev (nintedanib oral tablet)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

Treatment of idiopathic pulmonary fibrosis (IPF)

Treatment of chronic fibrosing interstitial lung diseases (ILDs) with a progressive phenotype

Slowing the rate of decline in pulmonary function in patients with systemic sclerosis-associated interstitial lung disease (SSc-ILD)

All other indications are considered experimental/investigational and are not a covered benefit.

II. CRITERIA FOR INITIAL APPROVAL

A. Treatment of IPF

An authorization may be granted for 6 months when the following criteria are met:

1. Prescriber is a pulmonologist or being prescribed in consultation with a pulmonologist.
2. Member is at least 18 years of age.
3. Documented diagnosis of IPF by all of the following criteria:
 - a. Exclusion of other known causes of interstitial lung disease (e.g., domestic and occupational environmental exposures, connective tissue disease, and drug toxicity), as documented by the following:
 - i. ICD-10 Code J84.112 (Idiopathic pulmonary fibrosis);
 - AND
 - b. One of the following:
 - i. In patients not subjected to surgical lung biopsy, the presence of a usual interstitial pneumonia (UIP) pattern on high-resolution computed tomography (HRCT) revealing IPF or probable IPF;
 - OR
 - ii. In patients subjected to a lung biopsy, both HRCT and surgical lung biopsy pattern reveal IPF or probable IPF
4. Member is not concurrently taking in combination with Esbriet (pirfenidone).
5. Baseline liver function tests have been performed.

B. Treatment of chronic fibrosing ILDs with a progressive phenotype

An authorization may be granted for 6 months when the following criteria are met:

1. Prescriber is a pulmonologist or being prescribed in consultation with a pulmonologist.
2. Member is at least 18 years of age.
3. Diagnosis of chronic ILDs with a progressive phenotype as documented by both of the following:
 - a. Presence of fibrotic ILD as determined by finding evidence of pulmonary fibrosis on HRCT, involving at least 10% of the lungs
 - b. Patient is presenting with clinical signs of progression as defined by one of the following in the previous 24 months:
 - i. Forced vital capacity (FVC) decline of greater than 10%; OR
 - ii. Two of the following:
 1. FVC decline of greater than or equal to 5%, but less than 10%
 2. Patient is experiencing worsening respiratory symptoms
 3. Patient is exhibiting increasing extent of fibrotic changes on chest imaging.
4. Member is not concurrently taking in combination with Esbriet (pirfenidone).
5. Baseline liver function tests have been performed.

C. Slowing the rate of decline in pulmonary function with SSc-ILD

An authorization may be granted for 6 months when the following criteria are met:

1. Prescriber is a pulmonologist or being prescribed in consultation with a pulmonologist.
2. Member is at least 18 years of age.
 - a. Diagnosis of SSc-ILD As documented by all of the following: One of the following:
 - i. Skin thickening of the fingers of both hands extending proximal to the metacarpophalangeal joints; OR
 - b. At least two of the following:
 - i. Skin thickening of the fingers (e.g., puffy fingers, sclerodactyly of the fingers)
 - ii. Fingertip lesions (e.g., digital tip ulcers, fingertip pitting scars)
 - iii. Telangiectasia
 - iv. Abnormal nailfold capillaries
 - v. Pulmonary arterial hypertension
 - vi. Raynaud's phenomenon
 - vii. SSc-related autoantibodies (e.g., anticentromere, antitopoisomerase I, anti-RNA polymerase III)
3. Member is not concurrently taking in combination with Esbriet (pirfenidone).
4. Baseline liver function tests have been performed.

III. CONTINUATION OF THERAPY

Authorization of 12 months may be granted when the member has documentation of a positive clinical response to the medication, is not being used in combination with Esbriet (pirfenidone), and is being prescribed by a pulmonologist or in consultation with a pulmonologist.

Effective Date: 12/01/2020
Last Reviewed: 9/2020
Scope: Medicaid

IV. QUANTITY LIMIT

Ofev has a quantity limit of #60 tablets per 30 days for the 100mg and 150mg.

V. REFERENCES

1. Ofev (nintedanib). Boehringer Ingelheim Pharmaceuticals. Ridgefield, CT. FDA Package Insert. March 2020.