

STEP THERAPY WITH QUANTITY LIMIT AND POST LIMIT PRIOR AUTHORIZATION CRITERIA

DRUG CLASS	EXTENDED-RELEASE OPIOID ANALGESICS
BRAND NAME (generic)	ARYMO ER (morphine sulfate extended-release tablets)
	AVINZA (morphine extended-release capsules)
	BELBUCA (buprenorphine buccal film)
	BUTRANS (buprenorphine transdermal system)
	CONZIP (tramadol hydrochloride extended-release capsules)
	DOLOPHINE 5 MG, 10 MG (methadone hydrochloride tablets)
	DURAGESIC (fentanyl transdermal system)
	EXALGO (hydromorphone hydrochloride extended-release tablets)
	HYSINGLA ER (hydrocodone bitartrate extended-release tablets)
	KADIAN (morphine extended-release capsules)
	METHADONE 5 MG, 10 MG (methadone hydrochloride tablets)
	METHADONE 200 MG/20 ML INJ (methadone hydrochloride injection)

METHADONE INTENSOL 10 MG/ML
(methadone oral concentrate)

METHADONE 5 MG/5 ML & 10 MG/5 ML ORAL SOLN
(methadone hydrochloride oral solution)

MORPHABOND ER
(morphine extended-release tablets)

MS CONTIN
(morphine extended-release tablets)

NUCYNTA ER
(tapentadol extended-release tablets)

OPANA ER
(oxymorphone hydrochloride extended-release tablets)

OXYCONTIN
(oxycodone hydrochloride extended-release tablets)

(oxymorphone hydrochloride extended-release tablets)

TARGINIQ ER
(oxycodone HCl/naloxone HCl extended-release tablets)

(tramadol hydrochloride extended-release)

TROXYCA ER
(oxycodone hydrochloride/naltrexone extended-release capsules)

ULTRAM ER
(tramadol hydrochloride extended-release tablets)

VANTRELA ER
(hydrocodone bitartrate extended-release tablets)

XTAMPZA ER
(oxycodone extended-release capsules)

ZOHYDRO ER
(hydrocodone bitartrate extended-release capsules)

Status: CVS Caremark Criteria

POLICY

FDA-APPROVED INDICATIONS

Arymo ER, Avinza, Kadian, MorphaBond ER, and MS Contin

Arymo ER, Avinza, Kadian, MorphaBond ER, and MS Contin are indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Arymo ER, Avinza, Kadian, MorphaBond ER, and MS Contin for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Arymo ER, Avinza, Kadian, MorphaBond ER, and MS Contin are not indicated as an as-needed (prn) analgesic.

Belbuca and Butrans

Belbuca and Butrans are indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioid formulations, reserve Belbuca and Butrans for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Belbuca and Butrans are not indicated as an as-needed (prn) analgesic.

ConZip, Ultram ER, and Tramadol Hydrochloride Extended-Release

ConZip, Ultram ER, and Tramadol Hydrochloride Extended-Release are indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release/long-acting opioid formulations, reserve ConZip, Ultram ER, and Tramadol Hydrochloride Extended-Release for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- ConZip, Ultram ER, and Tramadol Hydrochloride Extended-Release is not indicated as an as-needed (prn) analgesic.

Dolophine Tablets

Dolophine tablets are indicated for the:

- Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioids, reserve Dolophine tablets for use in patients for whom alternative analgesic treatment options (e.g., non-opioid analgesics or immediate-release opioid analgesics) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Dolophine tablets are not indicated as an as-needed (prn) analgesic.
- Detoxification treatment of opioid addiction (heroin or other morphine-like drugs).
- Maintenance treatment of opioid addiction (heroin or other morphine-like drugs), in conjunction with appropriate social and medical services.

Limitations of Use

- Methadone products used for the treatment of opioid addiction in detoxification or maintenance programs are subject to the conditions for distribution and use required under 42 CFR 8.12.

Duragesic

Duragesic is indicated for the management of pain in opioid-tolerant patients, severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Patients considered opioid-tolerant are those who are taking, for one week or longer, at least 60 mg morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, 60 mg hydrocodone per day, or an equianalgesic dose of another opioid.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release/long-acting opioid formulations, reserve Duragesic for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Duragesic is not indicated as an as-needed (prn) analgesic.

Exalgo

Exalgo is indicated for the management of pain in opioid-tolerant patients severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Patients considered opioid tolerant are those who are receiving, for one week or longer, at least 60 mg oral morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, 60 mg oral hydrocodone per day, or an equianalgesic dose of another opioid.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Exalgo for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Exalgo is not indicated as an as-needed (prn) analgesic.

Hysingla ER

Hysingla ER is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Hysingla ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Hysingla ER is not indicated as an as-needed (prn) analgesic.

Methadone Injection

Methadone injection is indicated:

- For the management of pain severe enough to require an opioid analgesic and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses reserve methadone hydrochloride injection for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or opioid combination products):
 - Have not been tolerated, or are not expected to be tolerated,
 - Have not provided adequate analgesia, or are not expected to provide adequate analgesia.
- For use in temporary treatment of opioid dependence in patients unable to take oral medication.

Limitations of Use

- Injectable methadone products are not approved for the outpatient treatment of opioid dependence. In this patient population, parenteral methadone is to be used only for patients unable to take oral medication, such as hospitalized patients.

Conditions For Distribution And Use Of Methadone Products For The Treatment Of Opioid Addiction

Code of Federal Regulations, Title 42, Sec 8

Methadone products when used for the treatment of opioid addiction in detoxification or maintenance programs, shall be dispensed only by opioid treatment programs (and agencies, practitioners or institutions by formal agreement with the program sponsor) certified by the Substance Abuse and Mental Health Services Administration and approved by the designated state authority. Certified treatment programs shall dispense and use methadone in oral form only and according to the treatment requirements stipulated in the Federal Opioid Treatment Standards (42 CFR 8.12). See below for important regulatory exceptions to the general requirement for certification to provide opioid agonist treatment. Failure to abide by the requirements in these regulations may result in criminal prosecution, seizure of the drug supply, revocation of the program approval, and injunction precluding operation of the program.

Regulatory Exceptions To The General Requirement For Certification To Provide Opioid Agonist Treatment:

During inpatient care, when the patient was admitted for any condition other than concurrent opioid addiction [pursuant to 21CFR 1306.07(c)], to facilitate the treatment of the primary admitting diagnosis.

During an emergency period of no longer than 3 days while definitive care for the addiction is being sought in an appropriately licensed facility [pursuant to 21CFR 1306.07(b)].

Methadone Intensol

Methadone Hydrochloride Intensol (oral concentrate) is indicated for the:

- Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioids, reserve Methadone Hydrochloride Intensol for use in patients for whom alternative analgesic treatment options (e.g., non-opioid analgesics or immediate-release opioid analgesics) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Methadone is not indicated as an as-needed (prn) analgesic.
- Detoxification treatment of opioid addiction (heroin or other morphine-like drugs).
- Maintenance treatment of opioid addiction (heroin or other morphine-like drugs), in conjunction with appropriate social and medical services.

Limitations of Use

- Methadone products used for the treatment of opioid addiction in detoxification or maintenance programs are subject to the conditions for distribution and use required under 42 CFR 8.2.

Conditions For Distribution And Use Of Methadone Products For The Treatment Of Opioid Addiction

Code of Federal Regulations, Title 42, Sec 8

Methadone products when used for the treatment of opioid addiction in detoxification or maintenance programs, shall be dispensed only by opioid treatment programs (and agencies, practitioners or institutions by formal agreement with the program sponsor) certified by the Substance Abuse and Mental Health Services Administration and approved by the designated state authority. Certified treatment programs shall dispense and use methadone in oral form only and according to the treatment requirements stipulated in the Federal Opioid Treatment Standards (42 CFR 8.12). See below for important regulatory exceptions to the general requirement for certification to provide opioid agonist treatment. Failure to abide by the requirements in these regulations may result in criminal prosecution, seizure of the drug supply, revocation of the program approval, and injunction precluding operation of the program.

Regulatory Exceptions To The General Requirement For Certification To Provide Opioid Agonist Treatment:

During inpatient care, when the patient was admitted for any condition other than concurrent opioid addiction [pursuant to 21CFR 1306.07(c)], to facilitate the treatment of the primary admitting diagnosis.

During an emergency period of no longer than 3 days while definitive care for the addiction is being sought in an appropriately licensed facility [pursuant to 21CFR 1306.07(b)].

Methadone Oral Solution

Methadone hydrochloride oral solution is indicated for the:

- Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioids, reserve methadone hydrochloride oral solution for use in patients for whom alternative analgesic treatment options (e.g., non-opioid analgesics or immediate-release

opioid analgesics) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.

- Methadone hydrochloride oral solution is not indicated as an as-needed (prn) analgesic.
- Detoxification treatment of opioid addiction (heroin or other morphine-like drugs).
- Maintenance treatment of opioid addiction (heroin or other morphine-like drugs), in conjunction with appropriate social and medical services.

Limitations of Use

- Methadone products used for the treatment of opioid addiction in detoxification or maintenance programs are subject to the conditions for distribution and use required under 42 CFR 8.2.

Methadone Tablets

Methadone hydrochloride tablets are indicated for the:

- Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with long-acting opioids, reserve methadone hydrochloride tablets for use in patients for whom alternative analgesic treatment options (e.g., non-opioid analgesics or immediate-release opioid analgesics) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Methadone hydrochloride tablets are not indicated as an as-needed (prn) analgesic.
- Detoxification treatment of opioid addiction (heroin or other morphine-like drugs).
- Maintenance treatment of opioid addiction (heroin or other morphine-like drugs), in conjunction with appropriate social and medical services.

Limitations of Use

- Methadone products used for the treatment of opioid addiction in detoxification or maintenance programs are subject to the conditions for distribution and use required under 42 CFR 8.2.

Nucynta ER

Nucynta ER is indicated for the management of:

- Pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.
- Neuropathic pain associated with diabetic peripheral neuropathy (DPN) in adults severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Nucynta ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Nucynta ER is not indicated as an as-needed (prn) analgesic.

Opana ER

Opana ER is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Opana ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Opana ER is not indicated as an as-needed (prn) analgesic.

OxyContin

OxyContin is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate in:

- Adults; and

- Opioid-tolerant pediatric patients 11 years of age and older who are already receiving and tolerate a minimum daily opioid dose of at least 20 mg oxycodone orally or its equivalent.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Oxycontin for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- OxyContin is not indicated as an as-needed (prn) analgesic.

Targiniq ER

Targiniq ER is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Targiniq ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Targiniq ER is not indicated as an as-needed (prn) analgesic.
- The maximum total daily dose of Targiniq ER should not exceed 80 mg/40 mg (40 mg/20 mg q12h) because higher doses may be associated with symptoms of opioid withdrawal or decreased analgesia.

Troxyca ER

Troxyca ER is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Troxyca ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Troxyca ER is not indicated as an as-needed (prn) analgesic.

Vantrela ER

Vantrela ER is an opioid agonist indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitation of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Vantrela ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Vantrela ER is not indicated as an as-needed (prn) analgesic.

Xtampza ER

Xtampza ER is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Xtampza ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Xtampza ER is not indicated as an as-needed (prn) analgesic.

Zohydro ER

Zohydro ER is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve Zohydro ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Zohydro ER is not indicated as an as-needed (prn) analgesic.

SCREENOUT LOGIC

If the patient has filled a prescription for at least a 1-day supply of a drug indicating the patient is being treated for cancer or sickle cell disease within the past 365 days under a prescription benefit administered by CVS Caremark, then the requested drug will be paid under that prescription benefit.

If a claim is submitted with an ICD 10 diagnosis code indicating cancer, sickle cell disease, or palliative care under a prescription benefit administered by CVS Caremark, then the requested drug will be paid under that prescription benefit.

If the patient has an ICD 10 diagnosis code indicating cancer or palliative care in their member health profile in the past 365 days, then the requested drug will be paid under that prescription benefit.

If the patient has any history of an ICD 10 diagnosis code indicating sickle cell disease in their member health profile, then the requested drug will be paid under that prescription benefit.

If a claim is submitted using a hospice patient residence code under a prescription benefit administered by CVS Caremark, then the requested drug will be paid under that prescription benefit.

INITIAL STEP THERAPY

For patients with no prescription claims of a cancer drug or a sickle cell disease drug in the past 365 days, no ICD 10 diagnosis code indicating cancer, sickle cell disease, or palliative care submitted with their prescription claim, no ICD 10 diagnosis code indicating cancer or palliative care in their member health profile in the past 365 days, no history of an ICD 10 diagnosis code indicating sickle cell disease in their member health profile, or no hospice patient residence code submitted with their prescription claim:

If the patient has filled a prescription for at least an 8-day supply of an immediate-release (IR) opioid agent indicated for the management of pain within prescription claim history in the past 90 days under a prescription benefit administered by CVS Caremark, then the initial quantity limit criteria will apply (see Column A and Column B in the Opioid Analgesics ER Quantity Limits Chart below).

If the patient has filled a prescription for at least a 30-day supply of an extended-release (ER) opioid agent indicated for the management of pain within prescription claim history in the past 90 days under a prescription benefit administered by CVS Caremark, then the initial quantity limit criteria will apply (see Column A and Column B in the Opioid Analgesics ER Quantity Limits Chart below).

If the patient does not have at least an 8-day supply of an IR opioid agent indicated for the management of pain OR at least a 30-day supply of an ER opioid agent indicated for the management of pain within prescription claim history in the past 90 days (i.e., the patient has not used an IR opioid prior to the ER opioid OR the patient is not already stable on an ER opioid), then the claim will reject with a message indicating that a prior authorization (PA) is required. The prior authorization criteria would then be applied to requests submitted for evaluation to the PA unit.

COVERAGE CRITERIA

The requested drug will be covered with prior authorization when the following criteria are met:

- The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through hospice or palliative care

OR

- The requested drug is being prescribed for CHRONIC pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid [Note: Chronic pain is generally defined as pain that typically lasts greater than 3 months.]
AND
- The patient can safely take the requested dose based on their history of opioid use
AND
- The patient has been evaluated and the patient will be monitored regularly for the development of opioid use disorder
AND
- The patient's pain will be reassessed in the first month after the initial prescription or any dose increase AND every 3 months thereafter to ensure that clinically meaningful improvement in pain and function outweigh risks to patient safety
AND
- This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days **OR**
- The patient has taken an immediate-release opioid for at least one week
AND
- If the request is for a methadone product, then it is NOT being prescribed for detoxification treatment or as part of a maintenance treatment plan for opioid/substance abuse or addiction

[Note: These drugs should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.]

Quantity Limits may apply.

Opioid Analgesics ER Quantity Limits Chart

Coverage is provided without prior authorization for a 30-day or 90-day supply of an extended-release opioid based on dosage frequency recommendations in product labeling (when Step Therapy criteria met). Coverage for quantities above the initial limit for a 30-day or 90-day supply is provided through prior authorization when coverage conditions are met.

These quantity limits should accumulate across all drugs of the same unit limit (i.e., drugs with 30 units accumulate together, drugs with 60 units accumulate together, etc).

		COLUMN A	COLUMN B	COLUMN C	COLUMN D
Drug/Strength	Recommended Dosing	Initial 1 Month Limit* (per 25 days)	Initial 3 Month Limit* (per 75 days)	Post 1 Month* Limit (per 25 days)	Post 3 Month* Limit (per 75 days)
Arymo ER 15 mg	q8-12h	90 tabs/month 3 tabs/day (45 MME/day)	270 tabs/3 months 3 tabs/day (45 MME/day)	120 tabs/month 4 tabs/day (60 MME/day)	360 tabs/3 months 4 tabs/day (60 MME/day)
Arymo ER 30 mg	q8-12h	90 tabs/month 3 tabs/day (90 MME/day)	270 tabs/3 months 3 tabs/day (90 MME/day)	120 tabs/month 4 tabs/day (120 MME/day)	360 tabs/3 months 4 tabs/day (120 MME/day)
Arymo ER 60 mg	q8-12h	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)	120 tabs/month 4 tabs/day (240 MME/day)	360 tabs/3 months 4 tabs/day (240 MME/day)
Avinza 30 mg	q24h, MAX 1600 mg/day	30 caps/month 1 cap/day (30 MME/day)	90 caps/3 months 1 cap/day (30 MME/day)	60 caps/month 2 caps/day (60 MME/day)	180 caps/3 months 2 caps/day (60 MME/day)
Avinza 45 mg	q24h, MAX 1600 mg/day	30 caps/month 1 cap/day (45 MME/day)	90 caps/3 months 1 cap/day (45 MME/day)	60 caps/month 2 caps/day (90 MME/day)	180 caps/3 months 2 caps/day (90 MME/day)
Avinza 60 mg	q24h, MAX 1600 mg/day	30 caps/month 1 cap/day	90 caps/3 months 1 cap/day	60 caps/month 2 caps/day	180 caps/3 months 2 caps/day

		(60 MME/day)	(60 MME/day)	(120 MME/day)	(120 MME/day)
Avinza 75 mg	q24h, MAX 1600 mg/day	30 caps/month 1 cap/day (75 MME/day)	90 caps/3 months 1 cap/day (75 MME/day)	60 caps/month 2 caps/day (150 MME/day)	180 caps/3 months 2 caps/day (150 MME/day)
Avinza 90 mg	q24h, MAX 1600 mg/day	30 caps/month 1 cap/day (90 MME/day)	90 caps/3 months 1 cap/day (90 MME/day)	60 caps/month 2 caps/day (180 MME/day)	180 caps/3 months 2 caps/day (180 MME/day)
Avinza 120 mg	q24h, MAX 1600 mg/day	30 caps/month 1 cap/day (120 MME/day)	90 caps/3 months 1 cap/day (120 MME/day)	60 caps/month 2 caps/day (240 MME/day)	180 caps/3 months 2 caps/day (240 MME/day)
Belbuca 75 mcg	q12h, MAX 900 mcg/12 hrs	60 films/month 2 films/day (4.5 MME/day)	180 films/3 months 2 films/day (4.5 MME/day)	90 films/month 3 films/day (6.75 MME/day)	270 films/3 months 3 films/day (6.75 MME/day)
Belbuca 150 mcg	q12h, MAX 900 mcg/12 hrs	60 films/month 2 films/day (9 MME/day)	180 films/3 months 2 films/day (9 MME/day)	90 films/month 3 films/day (13.5 MME/day)	270 films/3 months 3 films/day (13.5 MME/day)
Belbuca 300 mcg	q12h, MAX 900 mcg/12 hrs	60 films/month 2 films/day (18 MME/day)	180 films/3 months 2 films/day (18 MME/day)	90 films/month 3 films/day (27 MME/day)	270 films/3 months 3 films/day (27 MME/day)
Belbuca 450 mcg	q12h, MAX 900 mcg/12 hrs	60 films/month 2 films/day (27 MME/day)	180 films/3 months 2 films/day (27 MME/day)	90 films/month 3 films/day (40.5 MME/day)	270 films/3 months 3 films/day (40.5 MME/day)
Belbuca 600 mcg	q12h, MAX 900 mcg/12 hrs	60 films/month 2 films/day (36 MME/day)	180 films/3 months 2 films/day (36 MME/day)	90 films/month 3 films/day (54 MME/day)	270 films/3 months 3 films/day (54 MME/day)
Belbuca 750 mcg	q12h, MAX 900 mcg/12 hrs	60 films/month 2 films/day (45 MME/day)	180 films/3 months 2 films/day (45 MME/day)	90 films/month 3 films/day (67.5 MME/day)	270 films/3 months 3 films/day (67.5 MME/day)
Belbuca 900 mcg	q12h, MAX 900 mcg/12 hrs	60 films/month 2 films/day (54 MME/day)	180 films/3 months 2 films/day (54 MME/day)	90 films/month 3 films/day (81 MME/day)	270 films/3 months 3 films/day (81 MME/day)
Butrans 5 mcg/hr	q7d, MAX 20 mcg/hr	4 patches/month 0.144 patch/day (9 MME/day)	12 patches/3 months 0.144 patch/day (9 MME/day)	8 patches/month 0.287 patch/day (18 MME/day)	24 patches/3 months 0.287 patch/day (18 MME/day)
Butrans 7.5 mcg/hr	q7d, MAX 20 mcg/hr	4 patches/month 0.144 patch/day (13.5 MME/day)	12 patches/3 months 0.144 patch/day (13.5 MME/day)	8 patches/month 0.287 patch/day (27 MME/day)	24 patches/3 months 0.287 patch/day (27 MME/day)
Butrans 10 mcg/hr	q7d, MAX 20 mcg/hr	4 patches/month 0.144 patch/day (18 MME/day)	12 patches/3 months 0.144 patch/day (18 MME/day)	8 patches/month 0.287 patch/day (36 MME/day)	24 patches/3 months 0.287 patch/day (36 MME/day)
Butrans 15 mcg/hr	q7d, MAX 20 mcg/hr	4 patches/month 0.144 patch/day (27 MME/day)	12 patches/3 months 0.144 patch/day (27 MME/day)	8 patches/month 0.287 patch/day (54 MME/day)	24 patches/3 months 0.287 patch/day (54 MME/day)
Butrans 20 mcg/hr	q7d, MAX 20 mcg/hr	4 patches/month 0.144 patch/day (36 MME/day)	12 patches/3 months 0.144 patch/day (36 MME/day)	8 patches/month 0.287 patch/day (72 MME/day)	24 patches/3 months 0.287 patch/day (72 MME/day)
ConZip 100 mg	qd, MAX 300 mg/day	30 caps/month 1 cap/day (10 MME/day)	90 caps/3 months 1 cap/day (10 MME/day)	60 caps/month 2 caps/day (20 MME/day)	180 caps/3 months 2 caps/day (20 MME/day)
ConZip 200 mg	qd, MAX 300 mg/day	30 caps/month 1 cap/day (20 MME/day)	90 caps/3 months 1 cap/day (20 MME/day)	60 caps/month 2 caps/day (40 MME/day)	180 caps/3 months 2 caps/day (40 MME/day)
ConZip 300 mg	qd, MAX 300 mg/day	30 caps/month 1 cap/day (30 MME/day)	90 caps/3 months 1 cap/day (30 MME/day)	60 caps/month 2 caps/day (60 MME/day)	180 caps/3 months 2 caps/day (60 MME/day)
Dolophine 5 mg**	q8-12h	90 tabs/month 3 tabs/day	270 tabs/3 months 3 tabs/day	120 tabs/month 4 tabs/day	360 tabs/3 months 4 tabs/day

		(60 MME/day)	(60 MME/day)	(80 MME/day)	(80 MME/day)
Dolophine 10 mg**	q8-12h	90 tabs/month 3 tabs/day (240 MME/day)	270 tabs/3 months 3 tabs/day (240 MME/day)	120 tabs/month 4 tabs/day (320 MME/day)	360 tabs/3 months 4 tabs/day (320 MME/day)
Duragesic 12 mcg/hr	q72h	10 patches/month 0.334 patch/day (28.8 MME/day)	30 patches/3 months 0.334 patch/day (28.8 MME/day)	20 patches/month 0.667 patch/day (57.6 MME/day)	60 patches/3 months 0.667 patch/day (57.6 MME/day)
Duragesic 25 mcg/hr	q72h	10 patches/month 0.334 patch/day (60 MME/day)	30 patches/3 months 0.334 patch/day (60 MME/day)	20 patches/month 0.667 patch/day (120 MME/day)	60 patches/3 months 0.667 patch/day (120 MME/day)
Duragesic 37.5 mcg/hr	q72h	10 patches/month 0.334 patch/day (90 MME/day)	30 patches/3 months 0.334 patch/day (90 MME/day)	20 patches/month 0.667 patch/day (180 MME/day)	60 patches/3 months 0.667 patch/day (180 MME/day)
Duragesic 50 mcg/hr	q72h	10 patches/month 0.334 patch/day (120 MME/day)	30 patches/3 months 0.334 patch/day (120 MME/day)	20 patches/month 0.667 patch/day (240 MME/day)	60 patches/3 months 0.667 patch/day (240 MME/day)
Duragesic 62.5 mcg/hr	q72h	10 patches/month 0.334 patch/day (150 MME/day)	30 patches/3 months 0.334 patch/day (150 MME/day)	20 patches/month 0.667 patch/day (300 MME/day)	60 patches/3 months 0.667 patch/day (300 MME/day)
Duragesic 75 mcg/hr	q72h	10 patches/month 0.334 patch/day (180 MME/day)	30 patches/3 months 0.334 patch/day (180 MME/day)	20 patches/month 0.667 patch/day (360 MME/day)	60 patches/3 months 0.667 patch/day (360 MME/day)
Duragesic 87.5 mcg/hr	q72h	10 patches/month 0.334 patch/day (210 MME/day)	30 patches/3 months 0.334 patch/day (210 MME/day)	20 patches/month 0.667 patch/day (420 MME/day)	60 patches/3 months 0.667 patch/day (420 MME/day)
Duragesic 100 mcg/hr	q72h	10 patches/month 0.334 patch/day (240 MME/day)	30 patches/3 months 0.334 patch/day (240 MME/day)	20 patches/month 0.667 patch/day (480 MME/day)	60 patches/3 months 0.667 patch/day (480 MME/day)
Exalgo 8 mg	qd	30 tabs/month 1 tab/day (32 MME/day)	90 tabs/3 months 1 tab/day (32 MME/day)	60 tabs/month 2 tabs/day (64 MME/day)	180 tabs/3 months 2 tabs/day (64 MME/day)
Exalgo 12 mg	qd	30 tabs/month 1 tab/day (48 MME/day)	90 tabs/3 months 1 tab/day (48 MME/day)	60 tabs/month 2 tabs/day (96 MME/day)	180 tabs/3 months 2 tabs/day (96 MME/day)
Exalgo 16 mg	qd	30 tabs/month 1 tab/day (64 MME/day)	90 tabs/3 months 1 tab/day (64 MME/day)	60 tabs/month 2 tabs/day (128 MME/day)	180 tabs/3 months 2 tabs/day (128 MME/day)
Exalgo 32 mg	qd	30 tabs/month 1 tab/day (128 MME/day)	90 tabs/3 months 1 tab/day (128 MME/day)	60 tabs/month 2 tabs/day (256 MME/day)	180 tabs/3 months 2 tabs/day (256 MME/day)
Hysingla ER 20 mg	q24h	30 tabs/month 1 tab/day (20 MME/day)	90 tabs/3 months 1 tab/day (20 MME/day)	60 tabs/month 2 tabs/day (40 MME/day)	180 tabs/3 months 2 tabs/day (40 MME/day)
Hysingla ER 30 mg	q24h	30 tabs/month 1 tab/day (30 MME/day)	90 tabs/3 months 1 tab/day (30 MME/day)	60 tabs/month 2 tabs/day (60 MME/day)	180 tabs/3 months 2 tabs/day (60 MME/day)
Hysingla ER 40 mg	q24h	30 tabs/month 1 tab/day (40 MME/day)	90 tabs/3 months 1 tab/day (40 MME/day)	60 tabs/month 2 tabs/day (80 MME/day)	180 tabs/3 months 2 tabs/day (80 MME/day)
Hysingla ER 60 mg	q24h	30 tabs/month 1 tab/day (60 MME/day)	90 tabs/3 months 1 tab/day (60 MME/day)	60 tabs/month 2 tabs/day (120 MME/day)	180 tabs/3 months 2 tabs/day (120 MME/day)
Hysingla ER 80 mg	q24h	30 tabs/month 1 tab/day	90 tabs/3 months 1 tab/day	60 tabs/month 2 tabs/day	180 tabs/3 months 2 tabs/day

		(80 MME/day)	(80 MME/day)	(160 MME/day)	(160 MME/day)
Hysingla ER 100 mg	q24h	30 tabs/month 1 tab/day (100 MME/day)	90 tabs/3 months 1 tab/day (100 MME/day)	60 tabs/month 2 tabs/day (200 MME/day)	180 tabs/3 months 2 tabs/day (200 MME/day)
Hysingla ER 120 mg	q24h	30 tabs/month 1 tab/day (120 MME/day)	90 tabs/3 months 1 tab/day (120 MME/day)	60 tabs/month 2 tabs/day (240 MME/day)	180 tabs/3 months 2 tabs/day (240 MME/day)
Kadian 10 mg	q12-24h	60 caps/month 2 caps/day (20 MME/day)	180 caps/3 months 2 caps/day (20 MME/day)	90 caps/month 3 caps/day (30 MME/day)	270 caps/3 months 3 caps/day (30 MME/day)
Kadian 20 mg	q12-24h	60 caps/month 2 caps/day (40 MME/day)	180 caps/3 months 2 caps/day (40 MME/day)	90 caps/month 3 caps/day (60 MME/day)	270 caps/3 months 3 caps/day (60 MME/day)
Kadian 30 mg	q12-24h	60 caps/month 2 caps/day (60 MME/day)	180 caps/3 months 2 caps/day (60 MME/day)	90 caps/month 3 caps/day (90 MME/day)	270 caps/3 months 3 caps/day (90 MME/day)
Kadian 40 mg	q12-24h	60 caps/month 2 caps/day (80 MME/day)	180 caps/3 months 2 caps/day (80 MME/day)	90 caps/month 3 caps/day (120 MME/day)	270 caps/3 months 3 caps/day (120 MME/day)
Kadian 50 mg	q12-24h	60 caps/month 2 caps/day (100 MME/day)	180 caps/3 months 2 caps/day (100 MME/day)	90 caps/month 3 caps/day (150 MME/day)	270 caps/3 months 3 caps/day (150 MME/day)
Kadian 60 mg	q12-24h	60 caps/month 2 caps/day (120 MME/day)	180 caps/3 months 2 caps/day (120 MME/day)	90 caps/month 3 caps/day (180 MME/day)	270 caps/3 months 3 caps/day (180 MME/day)
Kadian 80 mg	q12-24h	60 caps/month 2 caps/day (160 MME/day)	180 caps/3 months 2 caps/day (160 MME/day)	90 caps/month 3 caps/day (240 MME/day)	270 caps/3 months 3 caps/day (240 MME/day)
Kadian 100 mg	q12-24h	60 caps/month 2 caps/day (200 MME/day)	180 caps/3 months 2 caps/day (200 MME/day)	90 caps/month 3 caps/day (300 MME/day)	270 caps/3 months 3 caps/day (300 MME/day)
Kadian 200 mg	q12-24h	60 caps/month 2 caps/day (400 MME/day)	180 caps/3 months 2 caps/day (400 MME/day)	90 caps/month 3 caps/day (600 MME/day)	270 caps/3 months 3 caps/day (600 MME/day)
Methadone 5 mg**	q8-12h	90 tabs/month 3 tabs/day (60 MME/day)	270 tabs/3 months 3 tabs/day (60 MME/day)	120 tabs/month 4 tabs/day (80 MME/day)	360 tabs/3 months 4 tabs/day (80 MME/day)
Methadone 10 mg**	q8-12h	90 tabs/month 3 tabs/day (240 MME/day)	270 tabs/3 months 3 tabs/day (240 MME/day)	120 tabs/month 4 tabs/day (320 MME/day)	360 tabs/3 months 4 tabs/day (320 MME/day)
Methadone 200 mg/20 mL injection**	q8-12h	20 mL/month (1 multidose vial) 0.667 mL/day (26.7 MME/day)	60 mL/3 months (3 multidose vials) 0.667 mL/day (26.7 MME/day)	40 mL/month (2 multidose vials) 1.334 mL/day (53.3 MME/day)	120 mL/3 months (6 multidose vials) 1.334 mL/day (53.3 MME/day)
Methadone 10 mg/mL Intensol soln**	q8-12h	90 mL/month 3 mL/day (240 MME/day)	270 mL/3 months 3 mL/day (240 MME/day)	120 mL/month 4 mL/day (320 MME/day)	360 mL/3 months 4 mL/day (320 MME/day)
Methadone 5 mg/5 mL Oral soln**	q8-12h	450 mL/month 15 mL/day (60 MME/day)	1350 mL/3 months 15 mL/day (60 MME/day)	600 mL/month 20 mL/day (80 MME/day)	1800 mL/month 20 mL/day (80 MME/day)
Methadone 10 mg/5 mL Oral soln**	q8-12h	450 mL/month 15 mL/day (240 MME/day)	1350 mL/3 months 15 mL/day (240 MME/day)	600 mL/month 20 mL/day (320 MME/day)	1800 mL/month 20 mL/day (320 MME/day)
MorphaBond ER 15 mg	q8-12h	90 tabs/month 3 tabs/day (45 MME/day)	270 tabs/3 months 3 tabs/day (45 MME/day)	120 tabs/month 4 tabs/day (60 MME/day)	360 tabs/3 months 4 tabs/day (60 MME/day)
MorphaBond ER 30 mg	q8-12h	90 tabs/month 3 tabs/day (90 MME/day)	270 tabs/3 months 3 tabs/day (90 MME/day)	120 tabs/month 4 tabs/day (120 MME/day)	360 tabs/3 months 4 tabs/day (120 MME/day)
MorphaBond ER 60 mg	q8-12h	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)	120 tabs/month 4 tabs/day (240 MME/day)	360 tabs/3 months 4 tabs/day (240 MME/day)

MorphaBond ER 100 mg	q8-12h	90 tabs/month 3 tabs/day (300 MME/day)	270 tabs/3 months 3 tabs/day (300 MME/day)	120 tabs/month 4 tabs/day (400 MME/day)	360 tabs/3 months 4 tabs/day (400 MME/day)
MS Contin 15 mg	q8-12h	90 tabs/month 3 tabs/day (45 MME/day)	270 tabs/3 months 3 tabs/day (45 MME/day)	120 tabs/month 4 tabs/day (60 MME/day)	360 tabs/3 months 4 tabs/day (60 MME/day)
MS Contin 30 mg	q8-12h	90 tabs/month 3 tabs/day (90 MME/day)	270 tabs/3 months 3 tabs/day (90 MME/day)	120 tabs/month 4 tabs/day (120 MME/day)	360 tabs/3 months 4 tabs/day (120 MME/day)
MS Contin 60 mg	q8-12h	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)	120 tabs/month 4 tabs/day (240 MME/day)	360 tabs/3 months 4 tabs/day (240 MME/day)
MS Contin 100 mg	q8-12h	90 tabs/month 3 tabs/day (300 MME/day)	270 tabs/3 months 3 tabs/day (300 MME/day)	120 tabs/month 4 tabs/day (400 MME/day)	360 tabs/3 months 4 tabs/day (400 MME/day)
MS Contin 200 mg	q8-12h	90 tabs/month 3 tabs/day (600 MME/day)	270 tabs/3 months 3 tabs/day (600 MME/day)	120 tabs/month 4 tabs/day (800 MME/day)	360 tabs/3 months 4 tabs/day (800 MME/day)
Nucynta ER 50 mg	q12h, MAX 500 mg/day	60 tabs/month 2 tabs/day (40 MME/day)	180 tabs/3 months 2 tabs/day (40 MME/day)	90 tabs/month 3 tabs/day (60 MME/day)	270 tabs/3 months 3 tabs/day (60 MME/day)
Nucynta ER 100 mg	q12h, MAX 500 mg/day	60 tabs/month 2 tabs/day (80 MME/day)	180 tabs/3 months 2 tabs/day (80 MME/day)	90 tabs/month 3 tabs/day (120 MME/day)	270 tabs/3 months 3 tabs/day (120 MME/day)
Nucynta ER 150 mg	q12h, MAX 500 mg/day	60 tabs/month 2 tabs/day (120 MME/day)	180 tabs/3 months 2 tabs/day (120 MME/day)	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)
Nucynta ER 200 mg	q12h, MAX 500 mg/day	60 tabs/month 2 tabs/day (160 MME/day)	180 tabs/3 months 2 tabs/day (160 MME/day)	90 tabs/month 3 tabs/day (240 MME/day)	270 tabs/3 months 3 tabs/day (240 MME/day)
Nucynta ER 250 mg	q12h, MAX 500 mg/day	60 tabs/month 2 tabs/day (200 MME/day)	180 tabs/3 months 2 tabs/day (200 MME/day)	90 tabs/month 3 tabs/day (300 MME/day)	270 tabs/3 months 3 tabs/day (300 MME/day)
Opana ER 5 mg	q12h	60 tabs/month 2 tabs/day (30 MME/day)	180 tabs/3 months 2 tabs/day (30 MME/day)	90 tabs/month 3 tabs/day (45 MME/day)	270 tabs/3 months 3 tabs/day (45 MME/day)
Opana ER 7.5 mg	q12h	60 tabs/month 2 tabs/day (45 MME/day)	180 tabs/3 months 2 tabs/day (45 MME/day)	90 tabs/month 3 tabs/day (67.5 MME/day)	270 tabs/3 months 3 tabs/day (67.5 MME/day)
Opana ER 10 mg	q12h	60 tabs/month 2 tabs/day (60 MME/day)	180 tabs/3 months 2 tabs/day (60 MME/day)	90 tabs/month 3 tabs/day (90 MME/day)	270 tabs/3 months 3 tabs/day (90 MME/day)
Opana ER 15 mg	q12h	60 tabs/month 2 tabs/day (90 MME/day)	180 tabs/3 months 2 tabs/day (90 MME/day)	90 tabs/month 3 tabs/day (135 MME/day)	270 tabs/3 months 3 tabs/day (135 MME/day)
Opana ER 20 mg	q12h	60 tabs/month 2 tabs/day (120 MME/day)	180 tabs/3 months 2 tabs/day (120 MME/day)	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)
Opana ER 30 mg	q12h	60 tabs/month 2 tabs/day (180 MME/day)	180 tabs/3 months 2 tabs/day (180 MME/day)	90 tabs/month 3 tabs/day (270 MME/day)	270 tabs/3 months 3 tabs/day (270 MME/day)
Opana ER 40 mg	q12h	60 tabs/month 2 tabs/day (240 MME/day)	180 tabs/3 months 2 tabs/day (240 MME/day)	90 tabs/month 3 tabs/day (360 MME/day)	270 tabs/3 months 3 tabs/day (360 MME/day)
OxyContin 10 mg	q12h	60 tabs/month 2 tabs/day (30 MME/day)	180 tabs/3 months 2 tabs/day (30 MME/day)	90 tabs/month 3 tabs/day (45 MME/day)	270 tabs/3 months 3 tabs/day (45 MME/day)
OxyContin 15 mg	q12h	60 tabs/month 2 tabs/day (45 MME/day)	180 tabs/3 months 2 tabs/day (45 MME/day)	90 tabs/month 3 tabs/day (67.5 MME/day)	270 tabs/3 months 3 tabs/day (67.5 MME/day)
OxyContin 20 mg	q12h	60 tabs/month 2 tabs/day	180 tabs/3 months 2 tabs/day	90 tabs/month 3 tabs/day	270 tabs/3 months 3 tabs/day

		(60 MME/day)	(60 MME/day)	(90 MME/day)	(90 MME/day)
OxyContin 30 mg	q12h	60 tabs/month 2 tabs/day (90 MME/day)	180 tabs/3 months 2 tabs/day (90 MME/day)	90 tabs/month 3 tabs/day (135 MME/day)	270 tabs/3 months 3 tabs/day (135 MME/day)
OxyContin 40 mg	q12h	60 tabs/month 2 tabs/day (120 MME/day)	180 tabs/3 months 2 tabs/day (120 MME/day)	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)
OxyContin 60 mg	q12h	60 tabs/month 2 tabs/day (180 MME/day)	180 tabs/3 months 2 tabs/day (180 MME/day)	90 tabs/month 3 tabs/day (270 MME/day)	270 tabs/3 months 3 tabs/day (270 MME/day)
OxyContin 80 mg	q12h	60 tabs/month 2 tabs/day (240 MME/day)	180 tabs/3 months 2 tabs/day (240 MME/day)	90 tabs/month 3 tabs/day (360 MME/day)	270 tabs/3 months 3 tabs/day (360 MME/day)
Targiniq ER 10 mg/5 mg	q12h, MAX 80 mg/40 mg (40 mg/20 mg q12h)	60 tabs/month 2 tabs/day (30 MME/day)	180 tabs/3 months 2 tabs/day (30 MME/day)	90 tabs/month 3 tabs/day (45 MME/day)	270 tabs/3 months 3 tabs/day (45 MME/day)
Targiniq ER 20 mg/10 mg	q12h, MAX 80 mg/40 mg (40 mg/20 mg q12h)	60 tabs/month 2 tabs/day (60 MME/day)	180 tabs/3 months 2 tabs/day (60 MME/day)	90 tabs/month 3 tabs/day (90 MME/day)	270 tabs/3 months 3 tabs/day (90 MME/day)
Targiniq ER 40 mg/20 mg	q12h, MAX 80 mg/40 mg (40 mg/20 mg q12h)	60 tabs/month 2 tabs/day (120 MME/day)	180 tabs/3 months 2 tabs/day (120 MME/day)	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)
Tramadol ER 100 mg	qd, MAX 300 mg/day	30 tabs/month 1 tab/day (10 MME/day)	90 tabs/3 months 1 tab/day (10 MME/day)	60 tabs/month 2 tabs/day (20 MME/day)	180 tabs/3 months 2 tabs/day (20 MME/day)
Tramadol ER 150 mg	qd, MAX 300 mg/day	30 caps 1 cap/day (15 MME/day)	90 caps/3 months 1 cap/day (15 MME/day)	60 caps/month 2 caps/day (30 MME/day)	180 caps/3 months 2 caps/day (30 MME/day)
Tramadol ER 200 mg	qd, MAX 300 mg/day	30 tabs/month 1 tab/day (20 MME/day)	90 tabs/3 months 1 tab/day (20 MME/day)	60 tabs/month 2 tabs/day (40 MME/day)	180 tabs/3 months 2 tabs/day (40 MME/day)
Tramadol ER 300 mg	qd, MAX 300 mg/day	30 tabs/month 1 tab/day (30 MME/day)	90 tabs/3 months 1 tab/day (30 MME/day)	60 tabs/month 2 tabs/day (60 MME/day)	180 tabs/3 months 2 tabs/day (60 MME/day)
Troxyca ER 10 mg/1.2 mg	q12h	60 caps/month 2 caps/day (30 MME/day)	180 caps/3 months 2 caps/day (30 MME/day)	90 caps/month 3 caps/day (45 MME/day)	270 caps/3 months 3 caps/day (45 MME/day)
Troxyca ER 20 mg/2.4 mg	q12h	60 caps/month 2 caps/day (60 MME/day)	180 caps/3 months 2 caps/day (60 MME/day)	90 caps/month 3 caps/day (90 MME/day)	270 caps/3 months 3 caps/day (90 MME/day)
Troxyca ER 30 mg/3.6 mg	q12h	60 caps/month 2 caps/day (90 MME/day)	180 caps/3 months 2 caps/day (90 MME/day)	90 caps/month 3 caps/day (135 MME/day)	270 caps/3 months 3 caps/day (135 MME/day)
Troxyca ER 40 mg/4.8 mg	q12h	60 caps/month 2 caps/day (120 MME/day)	180 caps/3 months 2 caps/day (120 MME/day)	90 caps/month 3 caps/day (180 MME/day)	270 caps/3 months 3 caps/day (180 MME/day)
Troxyca ER 60 mg/7.2 mg	q12h	60 caps/month 2 caps/day (180 MME/day)	180 caps/3 months 2 caps/day (180 MME/day)	90 caps/month 3 caps/day (270 MME/day)	270 caps/3 months 3 caps/day (270 MME/day)
Troxyca ER 80 mg/9.6 mg	q12h	60 caps/month 2 caps/day (240 MME/day)	180 caps/3 months 2 caps/day (240 MME/day)	90 caps/month 3 caps/day (360 MME/day)	270 caps/3 months 3 caps/day (360 MME/day)
Ultram ER 100 mg	qd, MAX 300 mg/day	30 tabs/month 1 tab/day (10 MME/day)	90 tabs/3 months 1 tab/day (10 MME/day)	60 tabs/month 2 tabs/day (20 MME/day)	180 tabs/3 months 2 tabs/day (20 MME/day)
Ultram ER 200 mg	qd, MAX 300 mg/day	30 tabs/month 1 tab/day (20 MME/day)	90 tabs/3 months 1 tab/day (20 MME/day)	60 tabs/month 2 tabs/day (40 MME/day)	180 tabs/3 months 2 tabs/day (40 MME/day)
Ultram ER 300 mg	qd, MAX 300 mg/day	30 tabs/month 1 tab/day (30 MME/day)	90 tabs/3 months 1 tab/day (30 MME/day)	60 tabs/month 2 tabs/day (60 MME/day)	180 tabs/3 months 2 tabs/day (60 MME/day)
Vantrela ER 15 mg	q12h, MAX 90	60 tabs/month	180 tabs/3 months	90 tabs/month	270 tabs/3 months

	mg q12h (180 mg/day)	2 tabs/day (30 MME/day)	2 tabs/day (30 MME/day)	3 tabs/day (45 MME/day)	3 tabs/day (45 MME/day)
Vantrela ER 30 mg	q12h, MAX 90 mg q12h (180 mg/day)	60 tabs/month 2 tabs/day (60 MME/day)	180 tabs/3 months 2 tabs/day (60 MME/day)	90 tabs/month 3 tabs/day (90 MME/day)	270 tabs/3 months 3 tabs/day (90 MME/day)
Vantrela ER 45 mg	q12h, MAX 90 mg q12h (180 mg/day)	60 tabs/month 2 tabs/day (90 MME/day)	180 tabs/3 months 2 tabs/day (90 MME/day)	90 tabs/month 3 tabs/day (135 MME/day)	270 tabs/3 months 3 tabs/day (135 MME/day)
Vantrela ER 60 mg	q12h, MAX 90 mg q12h (180 mg/day)	60 tabs/month 2 tabs/day (120 MME/day)	180 tabs/3 months 2 tabs/day (120 MME/day)	90 tabs/month 3 tabs/day (180 MME/day)	270 tabs/3 months 3 tabs/day (180 MME/day)
Vantrela ER 90 mg	q12h, MAX 90 mg q12h (180 mg/day)	60 tabs/month 2 tabs/day (180 MME/day)	180 tabs/3 months 2 tabs/day (180 MME/day)	90 tabs/month 3 tabs/day (270 MME/day)	270 tabs/3 months 3 tabs/day (270 MME/day)
Xtampza ER 9 mg	q12h, MAX 288 mg/day	60 caps/month 2 caps/day (30 MME/day)	180 caps/3 months 2 caps/day (30 MME/day)	90 caps/month 3 caps/day (45 MME/day)	270 caps/3 months 3 caps/day (45 MME/day)
Xtampza ER 13.5 mg	q12h, MAX 288 mg/day	60 caps/month 2 caps/day (45 MME/day)	180 caps/3 months 2 caps/day (45 MME/day)	90 caps/month 3 caps/day (67.5 MME/day)	270 caps/3 months 3 caps/day (67.5 MME/day)
Xtampza ER 18 mg	q12h, MAX 288 mg/day	60 caps/month 2 caps/day (60 MME/day)	180 caps/3 months 2 caps/day (60 MME/day)	120 caps/month 4 caps/day (120 MME/day)	360 caps/3 months 4 caps/day (120 MME/day)
Xtampza ER 27 mg	q12h, MAX 288 mg/day	60 caps/month 2 caps/day (90 MME/day)	180 caps/3 months 2 caps/day (90 MME/day)	180 caps/month 6 caps/day (270 MME/day)	540 caps/3 months 6 caps/day (270 MME/day)
Xtampza ER 36 mg	q12h, MAX 288 mg/day	60 caps/month 2 caps/day (120 MME/day)	180 caps/3 months 2 caps/day (120 MME/day)	240 caps/month 8 caps/day (480 MME/day)	720 caps/3 months 8 caps/day (480 MME/day)
Zohydro ER 10 mg	q12h	60 caps/month 2 caps/day (20 MME/day)	180 caps/3 months 2 caps/day (20 MME/day)	90 caps/month 3 caps/day (30 MME/day)	270 caps/3 months 3 caps/day (30 MME/day)
Zohydro ER 15 mg	q12h	60 caps/month 2 caps/day (30 MME/day)	180 caps/3 months 2 caps/day (30 MME/day)	90 caps/month 3 caps/day (45 MME/day)	270 caps/3 months 3 caps/day (45 MME/day)
Zohydro ER 20 mg	q12h	60 caps/month 2 caps/day (40 MME/day)	180 caps/3 months 2 caps/day (40 MME/day)	90 caps/month 3 caps/day (60 MME/day)	270 caps/3 months 3 caps/day (60 MME/day)
Zohydro ER 30 mg	q12h	60 caps/month 2 caps/day (60 MME/day)	180 caps/3 months 2 caps/day (60 MME/day)	90 caps/month 3 caps/day (90 MME/day)	270 caps/3 months 3 caps/day (90 MME/day)
Zohydro ER 40 mg	q12h	60 caps/month 2 caps/day (80 MME/day)	180 caps/3 months 2 caps/day (80 MME/day)	90 caps/month 3 caps/day (120 MME/day)	270 caps/3 months 3 caps/day (120 MME/day)
Zohydro ER 50 mg	q12h	60 caps/month 2 caps/day (100 MME/day)	180 caps/3 months 2 caps/day (100 MME/day)	90 caps/month 3 caps/day (150 MME/day)	270 caps/3 months 3 caps/day (150 MME/day)

*The duration of 25 days is used for a 30-day fill period and 75 days is used for a 90-day fill period to allow time for refill processing.

Limits are set up both as quantity versus time and daily dose edits.

**Calculating MME for methadone in clinical practice often involves a sliding-scale approach whereby the conversion factor increases with increasing dose.

REFERENCES

1. Arymo ER [package insert]. Wayne, PA: Zyla Life Sciences US Inc.; October 2019.
2. Avinza [package insert]. New York, NY: Pfizer, Inc.; April 2014.
3. Belbuca [package insert]. Raleigh, NC: BioDelivery Sciences International, Inc.; March 2021.
4. Butrans [package insert]. Stamford, CT: Purdue Pharma L.P.; March 2021.
5. ConZip [package insert]. Bridgewater, NJ: Vertical Pharmaceuticals, Inc.; September 2021.
6. Dolophine Tablets [package insert]. Eatontown, NJ: West-Ward Pharmaceuticals Corp.; June 2021.
7. Duragesic [package insert]. Titusville, NJ: Janssen Pharmaceuticals, Inc; March 2021.

8. Exalgo [package insert]. Webster Groves, MO: SpecGX LLC; June 2020.
9. Hysingla ER [package insert]. Stamford, CT: Purdue Pharma L.P.; March 2021.
10. Kadian [package insert]. Madison, NJ: Allergan USA, Inc.; October 2019.
11. Methadone Injection [package insert]. Morgantown, WV: Mylan Institutional LLC; June 2021.
12. Methadone Intensol Concentrate [package insert]. Eatontown, NJ: Hikma Pharmaceuticals USA Inc.; April 2021.
13. Methadone Oral Solution [package insert]. Eatontown, NJ: Hikma Pharmaceuticals USA Inc.; July 2020.
14. Methadone Tablets [package insert]. Mahwah, NJ: Glenmark Pharmaceuticals, Inc.; December 2019.
15. MorphaBond ER [package insert]. Basking Ridge, NJ: Daiichi Sankyo, Inc.; October 2019.
16. MS Contin [package insert]. Stamford, CT: Rhodes Pharmaceuticals L.P.; March 2021.
17. Nucynta ER [package insert]. Stoughton, MA: Collegium Pharmaceutical, Inc; March 2021.
18. Opana ER [package insert]. Malvern, PA: Endo Pharmaceuticals Inc.; March 2021.
19. OxyContin [package insert]. Stamford, CT: Purdue Pharma L.P.; October 2021.
20. Targiniq ER [package insert]. Stamford, CT: Purdue Pharma L.P.; September 2018.
21. Tramadol extended-release capsules [package insert]. Los Angeles, CA: SA3, LLC; September 2018.
22. Tramadol extended-release tablets [package insert]. Morgantown, WV: Mylan Pharmaceuticals Inc.; June 2021.
23. Troxyca ER [package insert]. New York, NY: Pfizer, Inc.; December 2016.
24. Ultram ER [package insert]. Titusville, NJ: Janssen Pharmaceuticals, Inc.; August 2017.
25. Vantrela ER [package insert]. North Wales, PA: Teva Pharmaceuticals USA, Inc.; January 2017.
26. Xtampza ER [package insert]. Cincinnati, OH: Patheon Pharmaceuticals; March 2021.
27. Zohydro ER [package insert]. Morristown, NJ: Currax Pharmaceuticals, LLC; March 2021.
28. Lexicomp Online, AHFS DI (Adult and Pediatric) Online. Hudson, OH: Wolters Kluwer Clinical Drug Information, Inc. <http://online.lexi.com/>. Accessed December 2021.
29. Micromedex (electronic version). Truven Health Analytics, Greenwood Village, Colorado, USA. <http://www.micromedexsolutions.com/>. Accessed December 2021.
30. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain — United States, 2016. *MMWR Recomm Rep*. 2016;65:1–49. Available at: <http://dx.doi.org/10.15585/mmwr.rr6501e1>. Accessed December 2021.
31. Palliative Care. NCCN Guidelines version 2.2021. Available at: http://www.nccn.org/professionals/physician_gls/pdf/palliative.pdf. Accessed December 2021.
32. Adult Cancer Pain. NCCN Guidelines version 2.2021. Available at: https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf. Accessed December 2021.
33. Chou R, Fanciullo G, Fine P, et al. Clinical Guidelines for the Use of Chronic Opioid Therapy in Chronic Noncancer Pain. *The Journal of Pain*. 2009;10:113-130.
34. National Heart, Lung, and Blood Institute. Evidence-Based Management of Sickle Cell Disease: Expert Panel Report, 2014. Available at: https://www.nhlbi.nih.gov/sites/default/files/media/docs/sickle-cell-disease-report%20020816_0.pdf. Accessed December 2021.