

SPECIALTY GUIDELINE MANAGEMENT

TALTZ (ixekizumab)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

- A. Moderate to severe plaque psoriasis in patients 6 years of age and older who are candidates for systemic therapy or phototherapy
- B. Adult patients with active psoriatic arthritis
- C. Adult patients with active ankylosing spondylitis
- D. Adult patients with active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation

All other indications are considered experimental/investigational and not medically necessary.

II. DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

- A. Plaque psoriasis (PsO)
 - 1. Initial requests:
 - i. Chart notes or medical record documentation of affected area(s) and body surface area (BSA) affected (if applicable).
 - ii. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 - 2. Continuation requests: Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.
- B. Ankylosing spondylitis (AS), axial spondyloarthritis (axSpA), and psoriatic arthritis (PsA)
 - 1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
 - 2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.

III. PRESCRIBER SPECIALTIES

This medication must be prescribed by or in consultation with one of the following:

- A. Plaque psoriasis: dermatologist

- B. Psoriatic arthritis: rheumatologist or dermatologist
- C. Ankylosing spondylitis and axial spondyloarthritis: rheumatologist

IV. CRITERIA FOR INITIAL APPROVAL

A. Plaque psoriasis (PsO)

1. Authorization of 12 months may be granted for members 6 years of age and older who have previously received a biologic or targeted synthetic drug (e.g., Sotyktu, Otezla) indicated for the treatment of moderate to severe plaque psoriasis.
2. Authorization of 12 months may be granted for members 6 years of age and older for the treatment of moderate to severe plaque psoriasis when any of the following criteria is met:
 - a. Crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
 - b. At least 10% of the body surface area (BSA) is affected.
 - c. At least 3% of body surface area (BSA) is affected and the member meets any of the following criteria:
 - i. Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin.
 - ii. Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine, and acitretin (see Appendix A).

B. Psoriatic arthritis (PsA)

1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Otezla) indicated for active psoriatic arthritis.
2. Authorization of 12 months may be granted for adult members for treatment of active psoriatic arthritis when either of the following criteria is met:
 - i. Member has mild to moderate disease and meets one of the following criteria:
 - a. Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.
 - b. Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix B), or another conventional synthetic drug (e.g., sulfasalazine).
 - c. Member has enthesitis or predominantly axial disease.
 - ii. Member has severe disease.

C. Ankylosing spondylitis (AS) and axial spondyloarthritis (axSpA)

1. Authorization of 12 months may be granted for adult members who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Xeljanz) indicated for active ankylosing spondylitis or active axial spondyloarthritis.
2. Authorization of 12 months may be granted for adult members for treatment of active ankylosing spondylitis or active axial spondyloarthritis when any of the following criteria is met:
 - i. Member has experienced an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
 - ii. Member has an intolerance or contraindication to two or more NSAIDs.

V. CONTINUATION OF THERAPY

A. Plaque psoriasis (PsO)

Authorization of 12 months may be granted for all members 6 years of age and older (including new members) who are using the requested medication for moderate to severe plaque psoriasis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:

1. Reduction in body surface area (BSA) affected from baseline
2. Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

B. Psoriatic arthritis (PsA)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for psoriatic arthritis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Axial disease
6. Skin and/or nail involvement

C. Ankylosing spondylitis (AS) and axial spondyloarthritis (axSpA)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for active ankylosing spondylitis or active axial spondyloarthritis and who achieve or maintain positive clinical response with the requested medication as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Functional status
2. Total spinal pain
3. Inflammation (e.g., morning stiffness)

VI. OTHER

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [PPD], an interferon-release assay [IGRA], or a chest x-ray)* within 6 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

* If the screening testing for TB is positive, there must be further testing to confirm there is no active disease. Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug.

VII. DOSAGE AND ADMINISTRATION

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

VIII. APPENDICES

Appendix A. Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine, or Acitretin

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Drug interaction
4. Cannot be used due to risk of treatment-related toxicity
5. Pregnancy or currently planning pregnancy
6. Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)

Appendix B: Examples of Contraindications to Methotrexate or Leflunomide

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia)
4. Elevated liver transaminases
5. History of intolerance or adverse event
6. Hypersensitivity
7. Interstitial pneumonitis or clinically significant pulmonary fibrosis
8. Myelodysplasia
9. Pregnancy or currently planning pregnancy
10. Renal impairment
11. Significant drug interaction

IX. REFERENCES

1. Taltz [package insert]. Indianapolis, IN: Eli Lilly and Company; July 2022.
2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol*. 2011;65(1):137-174.
3. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol*. 2009;61:451-485.
4. Griffiths CE, Reich K, Lebwohl M, et al. Comparison of ixekizumab with etanercept or placebo in moderate-to-severe psoriasis (UNCOVER-2 and UNCOVER-3): results from two phase 3 randomised trials. *Lancet*. 2015;386(9993):541-51.
5. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis*. 2020;79(6):700-712.
6. McInnes IB, Mease PJ, Kirkham B, et al. Secukinumab, a human anti-interleukin-17A monoclonal antibody, in patients with psoriatic arthritis (FUTURE 2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2015;386(9999):1137-46.
7. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019;80(4):1029-1072.
8. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on June 15, 2022 from: <https://www.cdc.gov/tb/topic/basics/risk.htm>.
9. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis* 2011;70:896-904.
10. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network

Recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Rheumatol.* 2019;71(10):1285-1299.

11. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheum.* 2018;71:5-32.
12. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis.* 2020;79(6):700-712.
13. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. *Nat Rev Rheumatol.* 2022;18(8):465-479.