SPECIALTY GUIDELINE MANAGEMENT

OTEZLA (apremilast)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

- 1. Moderate to severe plaque psoriasis
- 2. Active psoriatic arthritis

All other indications are considered experimental/investigational and are not a covered benefit.

II. CRITERIA FOR INITIAL APPROVAL

A. Moderate to severe plague psoriasis

- Authorization of 12 months may be granted for members who have previously received Otezla or any biologic disease-modifying antirheumatic drug (DMARD) indicated for the treatment of moderate to severe plaque psoriasis.
- 2. Authorization of 12 months may be granted for treatment of moderate to severe plaque psoriasis when all of the following criteria are met:
 - a. At least 5% of BSA is affected OR crucial body areas (i.e., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
 - b. Member meets any of the following criteria:
 - i. Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine or acitretin.
 - ii. Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine or acitretin (see Appendix A).

B. Active psoriatic arthritis (PsA)

Authorization of 12 months may be granted for treatment of active psoriatic arthritis (PsA).

III. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for all members (including new members) who meet all initial authorization criteria and achieve or maintain positive clinical response after at least 4 months of therapy with Otezla as evidenced by low disease activity or improvement in signs and symptoms of the condition.

IV. Appendix A: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine or Acitretin.

1. Alcoholism, alcoholic liver disease, or other chronic liver disease

© 2019 CVS Caremark. All rights reserved.

This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written permission from CVS Caremark. This document contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with CVS Caremark.



- 2. Breastfeeding
- 3. Drug interaction
- 4. Cannot be used due to risk of treatment-related toxicity
- 5. Pregnancy or planning pregnancy (male or female)
- 6. Significant comorbidity prohibits use of systemic agents (examples include liver or kidney disease, blood dyscrasias, uncontrolled hypertension)

V. REFERENCES

- 1. Otezla [package insert]. Summit, NJ: Celgene Corporation; June 2017.
- 2. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 4: Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol.* 2009;61:451-485.
- 3. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011;65(1):137-174.
- Coates LC, Kavanaugh A, Mease PJ, et al. Group for research and assessment of psoriasis and psoriatic arthritis 2015 treatment recommendation for psoriatic arthritis. *Arthritis Rheumatol.* 2016 May;68(5):1060-71.

© 2019 CVS Caremark. All rights reserved.



This document contains confidential and proprietary information of CVS Caremark and cannot be reproduced, distributed or printed without written